









Composition (%w/w)

Total Nitrogen	13.5%
Nitric Nitrogen	13.1%
Ammoniacal Nitrogen	0.4%
Calcium Oxide	16.5%
Magnesium Oxide	6%

Agronomical Targets



Compatibility

Compatible with other fertilizers, with the exception of Sulphate or Phosphate fertilizers. Therefore a separate tank is needed or fertilizers should be applied at different times.

Packaging













Van Iperen CalMag Horticultural Grade is a derivate of Calcium and Magnesium Nitrate. The product is recognizable by its unique flake shape which is easily soluble with record solubility levels.

Combining Calcium and Magnesium with Nitrates is useful to promote healthy vegetative growth, strong fruit setting and qualitative fruit growth. This is achieved thanks to healthy cell walls and high chlorophyll production.

Van Iperen CalMag Horticultural Grade is recommended from early flowering and until end of fruit growth stages.

- Improves leaf colour, fruit quality and shelf-life
- Developed for foliar application. Also suitable for fertigation in open field and greenhouses
- Soft for leaf tissue

Product Characteristics

- Low on heavy metals
- Low on Sodium and Chloride
- Easy to handle, dissolve and apply



Did you know?

Van Iperen offers the widest range of needs! Our Calcium Magnesium Derivate provide solutions for broadcasting, foliar

Let's make the green switch!

We are Van Iperen International a Dutch producer of Specialty Fertilizers and Biostimulants. We are eager to change the rules of the game in plant nutrition, by providing highly innovative solutions to growers for more sustainable agriculture. Your local Van Iperen Sales Manager will help you and guide you to make the green switch together.

www.vaniperen.com

Dosage | Fertigation

Crop	Application date	Min kg/ha/season	Max kg/ha/season	
Grapes	During vegetative growth During fruit filling	80	120	
Banana	During vegetative growth During fruit filling	150	250	
Citrus	During vegetative growth During fruit filling	100	200	
Olive	During vegetative growth During fruit filling	60	120	

Dosage | Foliar application

Crop	Application date	Min kg/ha/appl	Max kg/ha/appl	Conc. % (w/v)
Grapes	3 - 5 applications:As of fruit settingDuring vegetative growth	4	10	0.6 - 0.8
Arable crops	2 - 3 applications: • 10 - 14 days interval • As of vegetative growth	2	4	0.3 - 0.5
Banana	1 - 2 applications: • As of fruit setting • During vegetative growth	3	12	0.3 - 1.0
Citrus	1 - 2 applications:As of fruit settingDuring vegetative growth	3	12	0.3 - 1.0

In case of foliar feeding as part of a mix with crop protection products or other fertilizers, a compatibility test has to be done prior to preparing the spray-mix.

The mentioned indicated dosages and application stages are given as a guideline. Exact dosages, concentration and application stage are subject to local conditions, use of other fertilizers and can only be given after an objective diagnosis.

