



ALGAE SOLUTIONS

FoliaStim® B Liquid

Van Iperen FoliaStim® B Liquid is a pure and highly concentrated liquid Micronutrient fertilizer, formulated with 15% seaweed. Our product is a stable solution and is recommended as of bud development, prior to flowering. The high level of Boron in our formula improves cell division and cell wall formation. The presence of high quality seaweed supports the uptake and the transportation of nutrients and increases resistance against abiotic stress. Produced in our own plant Euroliquids.



Composition (%w/w)

Boron	2%
Seaweed extract	15%

Compatibility

Compatible with other fertilizers, with the exception of Calcium fertilizers. Therefore a separate tank is needed or fertilizers should be applied on different times. The pH of the tank solution should be above 4.

Packaging



- It overcomes stunted growth and increases drought and heat resistance.
- Easy to handle and apply
- Developed for foliar application
- Soft for leaf tissue

Product Characteristics

- Highly concentrated solution
- 15% w/w of high quality Canadian seaweed (*Ascophylum Nodosum*)
- Dark brown liquid
- Processing of seaweed according to unique method, guaranteeing optimal effect
- Production process certified according to ISO 9001:2015

Did you know?

Van Iperen offers a wide range of seaweed products! Our seaweed range is based on high quality Canadian "Ascophyllum Nodosum". Together with our pure seaweed products we provide solutions for foliar application, fertigation in open field and greenhouses and root-dipping prior to planting. The Van Iperen Seaweed range are part of the extended range of Iperen Specialty Fertilizers.









Where Nutrition meets Biostimulation™

Van Iperen International is a Dutch producer of Specialty Fertilizers. We offer you the widest range of fertilizers combining nutrition with biostimulation. For more information please contact your local Van Iperen Sales Manager or visit our website:

www.vaniperen.com

Grow for the best.

Dosage | Foliar application

Crop	Application date	Min L/ha/appl	Max L/ha/appl	Conc. % (w/v)
Rapeseed 	1 - 3 applications: • As of vegetative re-growth, C2 stage • 7 - 10 days interval	2	4	1.0 - 2.0
Sugarbeet 	1 - 3 applications: • As of 8 - 10 leaves stage • 7 - 10 days interval	2	5	1.0 - 2.0
Soybean 	1 - 3 applications: • As of 10 cm • Until flowering starts • 7 - 10 days interval	2	4	1.0 - 2.0
Sunflower 	1 - 3 applications: • As of 2 weeks after seeding • Until flower bud opening • 10 - 14 days interval	2	4	1.0 - 2.0
Cabbage 	1 - 3 applications: • As of transplanting • 10 - 14 days interval	2	3	1.0 - 2.0
Melon 	1 - 3 applications: • As of vegetative growth • Until flowering starts • 10 - 14 days interval	2	4	1.0 - 2.0
Vineyards 	1 - 3 applications: • As of 5 - 6 leaf stage • 10 - 14 days interval	1.5	3	1.0 - 2.0
Olive 	• During 15 days before flowering	2	3	1.0 - 2.0

In case of foliar feeding as part of a mix with crop protection products, a compatibility test has to be done prior to preparing the spray-mix.

The mentioned indicated dosages and application stages are given as a guideline. Exact dosages, concentration and application stage are subject to local conditions, use of other fertilizers and can only be given after an objective diagnosis.