

SPECIALS

Cal Boron Horticultural Grade

Van Iperen Cal Boron Horticultural Grade is a derivate of Calcium Nitrate and Boron fertilizer, which dissolves rapidly and completely. Our product is not sensitive to caking and is recommended as of early spring until fruit setting, especially as of bud development prior to flowering. The high level of Calcium and Boron in our formula improves the strength of cell walls and cell division. The presence of Nitrogen has a positive effect on the uptake of Calcium in the plant.



Composition (%w/w)

| | |
|---------------------|-------|
| Total Nitrogen | 15.1% |
| Nitric Nitrogen | 14.3% |
| Ammoniacal Nitrogen | 0.8% |
| Calcium Oxide | 27% |
| Boron | 0.2% |

Compatibility

Compatible with other fertilizers, with the exception of Sulphate or Phosphate fertilizers. Therefore a separate tank is needed or fertilizers should be applied at different times.

Packaging



- Improves fruit quality, stunted growth and shelf-life
- Easy to handle, dissolve and apply
- Developed for foliar application. Also suitable for fertigation in open field and greenhouses
- Soft for leaf tissue

Product Characteristics

- Low on heavy metals
- Low on Sodium and Chloride
- White flakes



Did you know?

Van Iperen offers the widest range of Calcium products, for all your Calcium needs! Our Calcium Magnesium Derivate range includes soluble and liquid products. Together with our Calcium Nitrates and our chelated Calcium we provide solutions for broadcasting, foliar application and fertigation in open field and high-tech soilless crops. The Van Iperen Calcium and Magnesium Derivates are part of the extended range of Iperen Specialty Fertilizers.






Where Nutrition meets Biostimulation™

Van Iperen International is a Dutch producer of Specialty Fertilizers. We offer you the widest range of fertilizers combining nutrition with biostimulation. For more information please contact your local Van Iperen Sales Manager or visit our website:






www.vaniperen.com

Grow for the best.

Dosage | Fertigation

| Crop | Application date | Conc. % (w/v) |
|--|--|---------------|
| Grapes  | <ul style="list-style-type: none"> During vegetative growth During fruit filling | 0.10 - 0.20 |
| Banana  | <ul style="list-style-type: none"> During vegetative growth During fruit filling | 0.03 - 0.15 |
| Citrus  | <ul style="list-style-type: none"> During vegetative growth During fruit filling | 0.03 - 0.15 |
| Olive  | <ul style="list-style-type: none"> During vegetative growth During fruit filling | 0.10 - 0.20 |
| Pip fruit  | <ul style="list-style-type: none"> During vegetative growth During fruit filling | 0.03 - 0.15 |

Dosage | Foliar application

| Crop | Application date | Min kg/ha/appl | Max kg/ha/appl | Conc. % (w/v) |
|---|---|----------------|----------------|---------------|
| Grapes  | 3 - 5 applications: <ul style="list-style-type: none"> As of fruit setting During vegetative growth | 4 | 10 | 0.6 - 0.8 |
| Arable crops  | 2 - 3 applications: <ul style="list-style-type: none"> 10 - 14 days interval As of vegetative growth | 2 | 4 | 0.3 - 0.5 |
| Banana  | 1 - 2 applications: <ul style="list-style-type: none"> As of fruit setting During vegetative growth | 3 | 12 | 0.3 - 1.0 |
| Citrus  | 1 - 2 applications: <ul style="list-style-type: none"> As of fruit setting During vegetative growth | 3 | 12 | 0.3 - 1.0 |
| Olive  | 3 - 5 applications: <ul style="list-style-type: none"> As of fruit setting During vegetative growth | 4 | 10 | 0.6 - 0.8 |

In case of foliar feeding as part of a mix with crop protection products, a compatibility test has to be done prior to preparing the spray-mix.

The mentioned indicated dosages and application stages are given as a guideline. Exact dosages, concentration and application stage are subject to local conditions, use of other fertilizers and can only be given after an objective diagnosis.