



NPK SOLUTIONS

Dubbi Liquid

6 - 3 - 6 + MgO + TE



Composition (%w/w)

Total Nitrogen	6%
Nitric Nitrogen	3.6%
Ammoniacal Nitrogen	2.4%
Phosphorus Pentoxide	3%
Potassium Oxide	6%
Magnesium Oxide	0.02%
Boron	0.0023%
Copper	0.0012%
Iron	0.016%
Manganese	0.001%
Molybdenum	0.0003%
Zinc	0.0008%

Compatibility

Compatible with other fertilizers, with the exception of Calcium fertilizers. Therefore a separate tank is needed or fertilizers should be applied on different times. The pH of the tank solution should be above 4.

Packaging



Van Iperen Dubbi Liquid | 6 - 3 - 6 + MgO + TE is a pure and concentrated liquid NPK with a high level of Nitrogen and Potassium combined with EDTA chelated micronutrients. Our product is a transparent, stable solution and is recommended at different phenological stages. The high level of Nitrogen and Potassium in our formula improves cell elongation and cell pressure. The presence of Magnesium and micro nutrients contributes to the production of chlorophyll and supports general plant metabolism. Produced in our own plant Euroliquid

- Improves vegetative growth and gives high quality buds and fruits
- Micronutrients 100% chelated and fully available
- Easy to handle and apply
- Developed for foliar application. Also suitable for fertigation in open field and greenhouses
- Soft for leaf tissue

Product Characteristics

- Low on heavy metals
- Low on Sodium and Chloride
- Production process certified according to ISO 9001:2015
- Dark brown liquid

Did you know?

Van Iperen offers a wide range of high quality Liquid NPKs in different ratios adapted to local needs. Our Liquid NPK range consists of pure and stable formulas based on food grade raw materials. The Van Iperen Liquid NPKs are part of the extended range of Iperen Specialty Fertilizers.





Where Nutrition meets Biostimulation™

Van Iperen International is a Dutch producer of Specialty Fertilizers. We offer you the widest range of fertilizers combining nutrition with biostimulation. For more information please contact your local Van Iperen Sales Manager or visit our website:







www.vaniperen.com

Grow for the best.

Dosage | Fertigation

Crop	Application date	Min L/ha/season	Max L/ha/season
Fruit trees 	<ul style="list-style-type: none"> • As of fruit setting • Until 4 weeks before harvesting 	150	400
Banana 	<ul style="list-style-type: none"> • During entire fertigation program 	200	500
Citrus 	<ul style="list-style-type: none"> • During entire fertigation program 	150	400
Potato 	<ul style="list-style-type: none"> • As of tuber setting • Until tuber ripening 	100	200

Dosage | Foliar application

Crop	Application date	Min L/ha/appl	Max L/ha/appl	Conc. % (w/v)
Arable crops 	3 applications: <ul style="list-style-type: none"> • During entire season • 10 - 14 days interval 	3	5	0.3 - 0.5
Banana 	2 - 4 applications: <ul style="list-style-type: none"> • Until flowering starts 	5	10	0.5 - 0.8
Citrus 	2 applications: <ul style="list-style-type: none"> • Until fruit setting 	5	8	0.5 - 1.0
Olive 	4 applications: <ul style="list-style-type: none"> • Spring application • During vegetative growth • During fruit setting • During fruit filling 	5	10	0.5 - 1.0
Potato 	3 applications: <ul style="list-style-type: none"> • As of vegetative growth • 10 - 14 days interval 	3	5	0.3 - 0.5
Stone fruit 	2 applications: <ul style="list-style-type: none"> • During vegetative growth • During fruit filling 	3	5	0.3 - 0.5

In case of foliar feeding as part of a mix with crop protection products, a compatibility test has to be done prior to preparing the spray-mix.

The mentioned indicated dosages and application stages are given as a guideline. Exact dosages, concentration and application stage are subject to local conditions, use of other fertilizers and can only be given after an objective diagnosis.